

Direct test of the MSW effect by the solar appearance term in beam experiments

Walter Winter^{a*}

^a*School of Natural Sciences, Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, USA*

Abstract

We discuss if one can verify the MSW effect in neutrino oscillations at a high confidence level in long-baseline experiments. We demonstrate that for long enough baselines at neutrino factories, the matter effect sensitivity is, as opposed to the mass hierarchy sensitivity, not suppressed by $\sin^2 2\theta_{13}$ because it is driven by the solar oscillations in the appearance probability. Furthermore, we show that for the parameter independent direct verification of the MSW effect at long-baseline experiments, a neutrino factory with a baseline of at least 6 000 km is needed. For superbeams, we do not find a 5σ discovery potential of the MSW effect independent of $\sin^2 2\theta_{13}$. We finally summarize different methods to test the MSW effect.

PACS: 14.60.Pq

Key words: Neutrino oscillations, Matter effects, MSW effect, long-baseline experiments

1. Introduction

It is now widely believed that neutrino oscillations are modified by matter effects, which is often referred to as the Mikheev-Smirnov-Wolfenstein (MSW) effect [1–3]. In this effect, the coherent forward scattering in matter by charged currents results in phase shifts in neutrino oscillations. Since ordinary matter consists of electrons, but no muons or taus, the W boson exchange causes a relative phase shift of the electron neutrino flavor. This relative phase shift then translates into changes of the neutrino oscillation probabilities.

The establishment of the LMA (Large Mixing Angle) solution in solar neutrino oscillations by the combined knowledge from SNO [4], KamLAND [5], and the other solar neutrino experiments has lead to “indirect” evidence for the MSW effect within the sun. A more direct test of these matter effects would be the “solar day-night effect” (see Ref. [6] and references therein), where the solar neutrino flux can (during the night) be enhanced through matter effects in the Earth due to regeneration effects [7]. So far, the solar day-night effect has not been discovered at a high con-

fidence level by Super-Kamiokande and SNO solar neutrino measurements [8,9]. A future very large water Cherenkov detector used for proton decay ($\sim 7 \times$ Super-Kamiokande) could establish this effect at the 4σ confidence level within ten years [10]. Similar tests could be performed with supernova neutrinos [11], which, however, have a strong (neutrino flux) model, detector position(s), and θ_{13} dependence [12]. In addition, strong matter effects can also occur in atmospheric neutrino oscillations in the Earth [13,14]. Since the muon neutrino disappearance probability is, to first order in $\alpha \equiv \Delta m_{21}^2/\Delta m_{31}^2$ and $\sin \theta_{13}$, not affected by Earth matter effects [15], testing the matter effects in atmospheric neutrinos is very difficult. However, the appearance signal of future long-baseline experiments is supposed to be very sensitive towards matter effects in atmospheric neutrino oscillations (see, for example, Refs. [16–20]). This makes the long-baseline test one natural candidate to directly discover the MSW effect at a very high confidence level.

Since the direct verification of the MSW effect would be another consistency check for our picture of neutrino oscillations, we study the potential of future long-baseline experiments to test

*E-mail: winter@ias.edu

matter versus vacuum oscillations. A similar measurement based upon matter effects in neutrino oscillations is the mass hierarchy sensitivity, which assumes that the matter effect is present and then tests the difference between the normal and inverted mass hierarchies. We will use this measurement in some cases for comparison in order to show the similarities and differences to the matter effect sensitivity. Note that the direct test of the MSW effect at neutrino factories was, for example, studied in Ref. [17]. Since at that time the parameter $\alpha \equiv \Delta m_{21}^2/\Delta m_{31}^2$ was very small for the LMA best-fit values, the contributions from the solar terms in the appearance probability were neglected and the MSW effect sensitivity was therefore determined to be strongly suppressed by $\sin^2 2\theta_{13}$. We will show that the now larger best-fit value of Δm_{21}^2 (and thus α) does not justify this assumption anymore.

2. Analytical motivation and qualitative discussion

For long-baseline beam experiments, the electron or muon neutrino appearance probability P_{app} (one of the probabilities $P_{e\mu}$, $P_{\mu e}$, $P_{e\bar{\mu}}$, $P_{\bar{\mu}e}$) is very sensitive to matter effects, whereas the disappearance probability $P_{\mu\mu}$ (or $P_{\bar{\mu}\bar{\mu}}$) is, to first order, not. The appearance probability can be expanded in the small hierarchy parameter $\alpha \equiv \Delta m_{21}^2/\Delta m_{31}^2$ and the small $\sin 2\theta_{13}$ up to the second order as [15, 21, 22]:

$$\begin{aligned} P_{\text{app}} &\simeq \sin^2 2\theta_{13} \sin^2 \theta_{23} \frac{\sin^2[(1 - \hat{A})\Delta]}{(1 - \hat{A})^2} \\ &\pm \alpha \sin 2\theta_{13} \sin \delta_{\text{CP}} \sin(\Delta) \xi(\hat{A}, \Delta) \\ &+ \alpha \sin 2\theta_{13} \cos \delta_{\text{CP}} \cos(\Delta) \xi(\hat{A}, \Delta) \\ &+ \alpha^2 \cos^2 \theta_{23} \sin^2 2\theta_{12} \frac{\sin^2(\hat{A}\Delta)}{\hat{A}^2}. \end{aligned} \quad (1)$$

Here $\Delta \equiv \Delta m_{31}^2 L/(4E)$, $\xi(\hat{A}, \Delta) = \sin 2\theta_{12} \cdot \sin 2\theta_{23} \cdot \sin(\hat{A}\Delta)/\hat{A} \cdot \sin[(1 - \hat{A})\Delta]/(1 - \hat{A})$, and $\hat{A} \equiv \pm(2\sqrt{2}G_F n_e E)/\Delta m_{31}^2$ with G_F the Fermi coupling constant and n_e the electron density in matter. The sign of the second term is positive for $\nu_e \rightarrow \nu_\mu$ or $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ and negative for $\nu_\mu \rightarrow \nu_e$ or $\bar{\nu}_e \rightarrow \bar{\nu}_\mu$. The sign of \hat{A} is determined by the

sign of Δm_{31}^2 and choosing neutrinos (plus) or antineutrinos (minus). Note that the matter effect in Eq. (1) enters via the matter potential \hat{A} , where the equation reduces to the vacuum case for $\hat{A} \rightarrow 0$ (*cf.*, Ref. [15]).

Since $\sin^2 2\theta_{13} > 0$ has not yet been established, any suppression by $\sin^2 2\theta_{13}$ would be a major disadvantage for a measurement. Therefore, let us first investigate the interesting limit $\sin^2 2\theta_{13} \rightarrow 0$. In this limit, only the fourth term in Eq. (1) survives, which is often referred to as the “solar term”, since the appearance signal in the limit $\theta_{13} \rightarrow 0$ corresponds to the contribution from the solar neutrino oscillations. It would vanish in the two-flavor limit (limit $\alpha \rightarrow 0$) and would grow proportional to $(\Delta m_{21}^2 L/(4E))^2$ in vacuum (limit $\hat{A} \rightarrow 0$), as one expects from the solar neutrino contribution in the atmospheric limit. Note that this term is equal for the normal and inverted mass hierarchies, which means that it cannot be used for the mass hierarchy sensitivity. In order to show its effect for the matter effect sensitivity compared to vacuum, we use $\Delta P \equiv P_{\text{app}}^{\text{matter}} - P_{\text{app}}^{\text{vac}}$. We find from Eq. (1)

$$\begin{aligned} \Delta P^{\theta_{13} \rightarrow 0} &\simeq \alpha^2 \cos^2 \theta_{23} \sin^2 2\theta_{12} \\ &\times \Delta^2 \left(\frac{\sin^2(\hat{A}\Delta)}{\hat{A}^2 \Delta^2} - 1 \right). \end{aligned} \quad (2)$$

Thus, this remaining effect does not depend on $\sin^2 2\theta_{13}$ and strongly increases with the baseline. In particular, the function $\sin^2(\hat{A}\Delta)/(\hat{A}^2 \Delta^2)$ is maximal (*i.e.*, unity) for $\hat{A}\Delta \rightarrow 0$ and has its first root for $\hat{A}\Delta = \pi$ at the “magic baseline” $L \sim 7500 \text{ km}$.² In the Earth, where Eq. (1) is valid because of the approximation $\Delta m_{21}^2 L/(4E) \ll 1$, we therefore have $\Delta P^{\theta_{13} \rightarrow 0} < 0$. This means that the matter effects will suppress the appearance probability, where maximal suppression is obtained at the magic baseline. For short baselines,

²At the magic baseline [23], the condition $\sin(\hat{A}\Delta) = 0$ makes all terms but the first in Eq. (1) disappear in order to allow a “clean” (degeneracy-free) measurement of $\sin^2 2\theta_{13}$. Note that the argument $\hat{A}\Delta$ evaluates to $\sqrt{2}/2 G_F n_e L$ independent of E and Δm_{31}^2 , which means that it only depends on the baseline L . This also implies that the MSW effect in the limit $\theta_{13} \rightarrow 0$ actually modifies the solar oscillation frequency, because the argument $\hat{A}\Delta$ in Eq. (2) does not depend on Δm_{31}^2 .

the expansion in Δ shows that $\Delta P^{\theta_{13} \rightarrow 0} \propto L^4$ strongly grows with the baseline, and for very long baselines, the bracket in Eq. (2) becomes close to -1 , which means that $\Delta P^{\theta_{13} \rightarrow 0} \propto L^2$ compensated the $1/L^2$ -dependence of the flux. Thus, we expect to be able to test the matter effect even for vanishing θ_{13} if the baseline is long enough.

There is, however, another important ingredient in these qualitative considerations: The statistics has to be good enough to detect the term suppressed by α^2 . For the current best-fit values, α^2 evaluates to $\sim 10^{-3}$. One can easily estimate that the statistics of superbeams will normally be too low to measure the solar term for this value of α^2 to a high accuracy: Let us compare the first and fourth terms in Eq. (1), which are suppressed by $\sin^2 2\theta_{13}$ and α^2 , respectively. If one assumes that the other factors in the first and fourth terms are of order unity (at least for $\Delta \sim \pi/2$ close to the first oscillation maximum), one can estimate for a specific experiment that the contribution from the α^2 -term only becomes significant if the $\sin^2 2\theta_{13}$ -sensitivity limit of this experiment is much better than α^2 . This condition is, in general, not satisfied for the proposed superbeams³ and could only be circumvented by a very long baseline, where the probability difference in Eq. (2) grows $\propto L^2$. For example, the NO ν A superbeam in the simulation of Ref. [24] would only lead to about four events with almost no dependence on the matter effect for $\theta_{13} \rightarrow 0$ (dominated by the intrinsic beam background). For neutrino factories, however, this order of α^2 should be accessible for long enough baselines. For example, for the neutrino factory NuFact-II of Ref. [25] at a baseline of 6000 km, we find for $\theta_{13} \rightarrow 0$ about 90 events in matter compared to 421 in vacuum.

Another interesting limit is the one of large values of $\sin^2 2\theta_{13}$, where the first term in Eq. (1) dominates, *i.e.*, for $\sin 2\theta_{13} \gg \alpha$, which is equivalent to $\sin^2 2\theta_{13} \gg 10^{-3}$. It is strongly enhanced close to the matter resonance, where the reso-

nance condition is given by $\hat{A} \rightarrow +1$. This condition evaluates to a resonance energy of ~ 9 GeV (for $\rho = 3.5$ g/cm³ and $\Delta m_{31}^2 = 2.5 \cdot 10^{-3}$ eV²) in the Earth's mantle, which is usually covered by a neutrino factory energy spectrum. Therefore, neutrino factories are supposed to be very sensitive to matter effects, which are in this limit driven by the atmospheric Δm_{31}^2 . Since the sign of \hat{A} depends on the mass hierarchy (and using neutrino or antineutrinos), it leads to a strong enhancement (+) or suppression (-) of the appearance probability. One therefore expects a very good sensitivity to the mass hierarchy for large enough $\sin^2 2\theta_{13}$ and long enough baselines. Similarly, one would expect a very good sensitivity to the matter effect itself compared to the vacuum case, since the appearance probability in matter becomes for long baselines very different from the vacuum case [17]. However, at least in the limit of small Δ or \hat{A} , the difference between the matter and vacuum probabilities is by about a factor of two smaller than the one between the normal and inverted hierarchy matter probabilities, since the vacuum probability lies in between the other two. One can easily understand this in terms of the matter potential which “pulls” the probabilities in two different directions apart from the vacuum case. In addition, it is well known that the correlation with δ_{CP} highly affects the mass hierarchy sensitivity in large regions of the parameters space (see, *e.g.*, Refs. [25, 26]). Similarly, one can expect this correlation will destroy the matter effect sensitivity, too. Therefore, for large values of $\sin^2 2\theta_{13}$, it is natural to assume that the test of the matter effect will be harder than the one of the mass hierarchy.

3. Analysis methods and experiment simulation

In general, we use a three-flavor analysis of neutrino oscillations, where we take into account statistics, systematics, correlations, and degeneracies [26–29]. The analysis is performed with the $\Delta\chi^2$ method using the GLoBES software [30].

For the sensitivity to the matter effect, we test the hypothesis of vacuum oscillations, *i.e.*, we compute the simulated event rates for vacuum

³In fact, for superbeams, the background from the intrinsic (beam) electron neutrinos limits the performance, which means that increasing the luminosity would not solve this problem.

and a normal mass hierarchy. Note that there is not a large dependence on the mass hierarchy in vacuum, though the event rates depend (even in vacuum) somewhat on the mass hierarchy by the third term in Eq. (1) (if one is far enough off the oscillation maximum). We then test this hypothesis of vacuum oscillations by switching on the (constant) matter density profile and fit the rates to the simulated ones using the $\Delta\chi^2$ method. In order to take into account correlations, we marginalize over all the oscillation parameters and test both the normal and inverted hierarchies. As a result, we obtain the minimum $\Delta\chi^2$ for the given set of true oscillation parameters which best fit the vacuum case.

For the mass hierarchy sensitivity, we compute the simulated rate vector for the chosen mass hierarchy and fit it with the opposite sign of Δm_{31}^2 . Thus, it is determined by the minimum $\Delta\chi^2$ at the $\text{sgn}(\Delta m_{31}^2)$ -degeneracy [26]. Note that for neutrino factories this minimum at the opposite sign of Δm_{31}^2 might be very difficult to find because of mixed degeneracies. Since we assume maximal mixing, the only relevant mixed degeneracy here is the $(\delta_{\text{CP}}, \theta_{13})$ -degeneracy [27] for the inverted Δm_{31}^2 . The correlations originate in the minimization of the six-dimensional fit manifold at the position of the $\text{sgn}(\Delta m_{31}^2)$ -degeneracy, *i.e.*, any solution with the opposite sign of Δm_{31}^2 fitting the original solution destroys the mass hierarchy sensitivity [25]. In addition, we assume a constant matter density profile with 5% uncertainty, which takes into account matter density uncertainties as well as matter profile effects [31–33].

For all measurements, we assume that each experiment will provide the best measurement of the leading atmospheric oscillation parameters at that time, *i.e.*, we use the information from the disappearance channels simultaneously. However, we have tested for this study that the disappearance channels do not significantly contribute to the matter effect sensitivity.⁴ Furthermore, for

the leading solar parameters, we take into account that the ongoing KamLAND experiment will improve the errors down to a level of about 10% on each Δm_{21}^2 and $\sin 2\theta_{12}$ [34, 35].

As experiments, we will mainly use neutrino factories based upon the representative **NuFact-II** from Ref. [25]. In its standard configuration, it uses muons with an energy of 50 GeV, 4 MW target power ($5.3 \cdot 10^{20}$ useful muon decays per year), a baseline of 3 000 km, and a magnetized iron detector with a fiducial mass of 50 kt. We choose a symmetric operation with 4 yr in each polarity. For the oscillation parameters, we use, if not stated otherwise, the current best-fit values $\Delta m_{31}^2 = 2.5 \cdot 10^{-3} \text{ eV}^2$, $\sin^2 2\theta_{23} = 1$, $\Delta m_{21}^2 = 8.2 \cdot 10^{-5} \text{ eV}^2$, and $\sin^2 2\theta_{12} = 0.83$ [36–39]. We only allow values for $\sin^2 2\theta_{13}$ below the CHOOZ bound $\sin^2 2\theta_{13} \lesssim 0.1$ [40] and do not make any special assumptions about δ_{CP} . However, we will show in some cases the results for chosen selected values of δ_{CP} .

4. Quantitative results

We show in Figure 1 the sensitivity to the MSW effect for **NuFact-II** as function of the true values of $\sin^2 2\theta_{13}$ and the baseline L , where $\delta_{\text{CP}} = 0$ and a normal mass hierarchy are assumed. The sensitivity is given above the curves at the shown confidence levels. Obviously, the experiment can verify the MSW effect for long enough baselines even for $\sin^2 2\theta_{13} = 0$. The vertical dashed line separates the region where this measurement is dominated by the first term (θ_{13} -dominated) and the fourth term (solar-dominated) in Eq. (1). It is drawn for $\sin^2 2\theta_{13} = 10^{-3} \sim \alpha^2$, *i.e.*, in this region all the terms of Eq. (1) have similar magnitudes. Obviously, the performance in the θ_{13} -dominated (atmospheric oscillation-dominated) regime is much better than the one in the solar-dominated regime, because the θ_{13} -terms provide information on the matter effects in addition to the solar term. In this figure, the curves are shown for different selected confidence levels. However, in order to really establish the effect, a minimum 5σ signal will be necessary. Therefore, we will only use the 5σ curves below.

In order to discuss the most relevant param-

⁴In fact, the disappearance channels alone could resolve the matter effects for very large L and large $\sin^2 2\theta_{13}$. However, in this region, the relative contribution of the disappearance $\Delta\chi^2$ to the total one is only at the percent level.




Figure 1. Sensitivity to the MSW effect for NuFact-II as function of the true value of $\sin^2 2\theta_{13}$ and the baseline L . For the simulated oscillation parameters, the current best-fit values, $\delta_{CP} = 0$, and a normal mass hierarchy are assumed, whereas the fit parameters are marginalized. Sensitivity is given at the shown confidence level on the upper sides of the curves.

ter dependencies and to compare the matter effect and mass hierarchy sensitivities, we show in Figure 2 these sensitivities for two different values of δ_{CP} . As we have tested, the true value of δ_{CP} is one of the major impact factors for these measurements. In addition, the mass hierarchy sensitivity is modified by a similar amount for a simulated inverted instead of normal mass hierarchy, whereas the matter effect sensitivity does not show this dependence (because the reference rate vector is computed for vacuum). As far as the dependence on Δm_{21}^2 is concerned, we have not found any significant dependence of the MSW effect sensitivity within the current allowed 3σ range $7.4 \cdot 10^{-5} \text{ eV}^2 \lesssim \Delta m_{21}^2 \lesssim 9.2 \cdot 10^{-5} \text{ eV}^2$ [37]. Hence, we show in Figure 2 the selected two values of δ_{CP} for estimates of the (true) parameter dependencies, since there are no major qualitative differences.

As one can see from this figure, the behavior of the MSW sensitivity for short baselines and large $\sin^2 2\theta_{13}$ is qualitatively similar to the one of the mass hierarchy sensitivity, because both measurements are dominated by the θ_{13} -terms of Eq. (1). However, as we have indicated in Section 2, the difference between the normal and inverted hierarchy matter rates is about a factor of two larger than the one between vacuum and matter rates (for any mass hierarchy). Thus, for large $\sin^2 2\theta_{13}$, the mass hierarchy sensitivity is better than the MSW sensitivity (better means that it works for shorter baselines). Note that the solar (fourth) term in Eq. (1) is not dependent on the mass hierarchy, which means that there is no mass hierarchy sensitivity for small values of $\sin^2 2\theta_{13}$. In general, there are three regions for the MSW effect sensitivity in Figure 2:

$\sin^2 2\theta_{13} \lesssim 10^{-5}$: Only the solar term in Eq. (1) is present. The MSW effect sensitivity therefore does not depend on δ_{CP} or the mass hierarchy.

$\sin^2 2\theta_{13} \gtrsim 10^{-2}$: The measurement is dominated by the first term in Eq. (1) with some contribution of the second and third terms, which means that there is some dependence on δ_{CP} .

$10^{-5} \lesssim \sin^2 2\theta_{13} \lesssim 10^{-2}$: In the intermediate region, these two effects are competing, which leads to the “bump” in the right panel of Figure 2. In particular, the relative contribution of the CP terms in Eq. (1) is quite large, which means that one expects the strongest δ_{CP} -dependence there.

For the MSW effect sensitivity, one can easily see from both panels of Figure 2 that for $\sin^2 2\theta_{13} \gtrsim 0.05$ a baseline of 3000 km would be sufficient, because in this case the θ_{13} -signal is strong enough to provide information on the matter effects. However, in this case, $\sin^2 2\theta_{13}$ will be discovered by a superbeam and it is unlikely that a neutrino factory will be built. For smaller values $\theta_{13} < 0.01$, longer baselines will be necessary. In particular, to have sensitivity to the matter effect independent of the true parameter




Figure 2. The sensitivity to the MSW effect (black curves) and to the mass hierarchy (gray curves) for NuFact-II as function of the true value of $\sin^2 2\theta_{13}$ and the baseline L (5σ only). For the simulated oscillation parameters, the current best-fit values, $\delta_{CP} = 0$ (left) or $\delta_{CP} = \pi/2$ (right), and a normal mass hierarchy are assumed, whereas the fit parameters are marginalized over (solid curves). Sensitivity to the respective quantity is given on the upper/right side of the curves. The dashed curves correspond to the MSW effect sensitivity without correlations, *i.e.*, for all the fit parameters fixed.

values, a neutrino factory baseline $L \gtrsim 6000$ km is a prerequisite. Therefore, this matter effect test is another nice argument for at least one very long neutrino factory baseline. Note that one can read off the impact of correlations with the oscillation parameters from the comparison between the dashed and solid black curves in Figure 2. If one just fixed all the oscillation parameters, one would obtain the dashed curves. In this case, one could come to the conclusion that a shorter baseline would be sufficient, which is not true for the complete marginalized analysis.

As we have discussed in Section 2, the MSW test is very difficult for superbeams. For the combination of T2K, NO ν A, and Reactor-II from Ref. [24], it is not even possible at the 90% confidence level for $\sin^2 2\theta_{13} = 0.1$ at the CHOOZ bound. However, for a very large superbeam upgrade at very long baselines, there would indeed be some sensitivity to the matter effect even for vanishing θ_{13} . For example, if one used the

T2HK setup from Ref. [25] and (hypothetically) put the detector to a longer baseline, one would have some matter effect sensitivity at the 3σ confidence level for selected baselines $L \gtrsim 5500$ km. For the “magic baseline” $L \sim 7500$ km, one could even have a 4σ signal, but 5σ would hardly be possible.

5. Summary and discussion

We have investigated the potential of long-baseline experiments to test the matter effect (MSW effect) in neutrino oscillations. In particular, we have discussed under what conditions one can directly verify this MSW effect compared to vacuum oscillations at a high confidence level.

Though it is generally known that beam experiments are, for sufficiently long baselines, very sensitive to matter effects, we have demonstrated that the θ_{13} -terms in the appearance signal have much less matter effect sensitivity than one may expect. Especially, the comparison with another

Source/Method (where tested)	θ_{13} -suppr.	Reach [Ref.]	Comments/Assumptions
Solar ν /Sun	No	6σ [41]	MSW effect in sun; by comparison between vacuum and matter (existing solar ν experiments)
Solar ν /Earth (“day-night”)	No	4σ [10]	By large Water Cherenkov detector used for proton decay
SN ν /Earth, one detector	No	n/a [12]	Observation as “dips” in spectrum, but no observation guaranteed (because of flux uncertainties); effects depend on $\sin^2 2\theta_{13}$; HyperK-like detector needed
SN ν /Earth, two detectors	No	$4\sigma - 5\sigma$ [11]	For SN distance 10 Kpc, $E_B = 3 \cdot 10^{53}$ ergs; at least two Super-K size detectors, depends on their positions
Atmospheric ν /Earth	Yes	4σ [42]	Estimate for 100 kt magn. iron detector computed for $\sin^2 2\theta_{13} = 0.1$
Superbeam/Earth $L \lesssim 5\,500$ km	Yes	2σ	Estimate for T2HK-like setup for $\sin^2 2\theta_{13} \gtrsim 0.05$ at $L = 3\,000$ km; strongly depends on $\sin^2 2\theta_{13}$ and δ_{CP}
Superbeam/Earth $L \gtrsim 5\,500$ km	No	$\sim 3\sigma - 4\sigma$	Estimate for T2HK-like setup independent of $\sin^2 2\theta_{13}$
ν -factory/Earth $L \lesssim 6\,000$ km	Yes	5σ	Reach for $\sin^2 2\theta_{13} \gtrsim 0.05$ at $L = 3\,000$ km ($\delta_{CP} = \pi/2$); strongly depends on $\sin^2 2\theta_{13}$ and δ_{CP}
ν -factory/Earth $L \gtrsim 6\,000$ km	No	$5\sigma - 8\sigma$	Range depending on δ_{CP} for $L = 6\,000$ km; for $L \gg 6\,000$ km much better reach, such as $\sim 12\sigma$ for $L = 7\,500$ km

Table 1

Different methods to test the MSW effect: Source and method (in which medium the MSW effect is tested), the suppression of the effect by θ_{13} , the potential confidence level reach (including reference, where applicable), and comments/assumptions which have led to this estimate.

matter effect-dominated measurement, *i.e.*, the mass hierarchy sensitivity, has shown that the MSW effect sensitivity is much weaker for short baselines $L \lesssim 5\,000$ km. Note that both of these measurements suffer from correlations and degeneracies especially for intermediate $\sin^2 2\theta_{13}$.

However, for long enough baselines $L \gtrsim 6\,000$ km and good enough statistics, the solar term in the appearance probability is sensitive to matter effects compared to vacuum, which means that the MSW effect sensitivity is not suppressed by $\sin^2 2\theta_{13}$ anymore. Note that the solar term is not sensitive to the mass hierarchy at all, but it is reduced in matter compared to vacuum. In summary, we have demonstrated that a neutrino

factory with a sufficiently long baseline would have good enough statistics for a 5σ MSW effect discovery independent of $\sin^2 2\theta_{13}$, where the solar term becomes indeed statistically accessible. However, a very long baseline superbeam upgrade, such as a T2HK-like experiment at the “magic baseline” $L \sim 7\,500$ km, could have some sensitivity to the solar appearance term at the 4σ confidence level.

This result has three major implications: First, it is another argument for at least one very long neutrino factory baseline, where the other purposes of such a baseline could be a “clean” (correlation- and degeneracy-free) $\sin^2 2\theta_{13}$ -measurement at the “magic base-

line” [23] and a very good mass hierarchy sensitivity for large enough $\sin^2 2\theta_{13}$. The verification of the MSW effect would be a little “extra” for such a baseline. In addition, note that the mass hierarchy sensitivity assumes that the matter effects are present, which means that some more evidence for the MSW effect would increase the consistency of this picture.

Second, the absence of the $\sin^2 2\theta_{13}$ -suppression in the solar appearance term means that the direct MSW test at a beam experiment could be competitive with others methods, for a summary, see Table 1. However, it could be also partly complementary: If $\sin^2 2\theta_{13}$ turned out to be large, it is the atmospheric oscillation frequency which would be modified by matter effects and not the solar one. In addition, the MSW effect in Earth matter could be a more “direct” test under controllable conditions, because the Earth’s mantle has been extensively studied by seismic wave geophysics. Note that for atmospheric neutrinos, this test is much harder, an example can be found in Ref. [42].

Third, we have demonstrated that the solar term in the appearance probability can really provide statistically significant information, which may also be useful for other applications. For example, the dependence of the solar appearance term on $\cos \theta_{23}$ instead of $\sin^2 2\theta_{23}$ in the disappearance probability could, for properly chosen baselines, be useful to resolve the $(\theta_{23}, \pi/2 - \theta_{23})$ -degeneracy [43]. Thus, the formerly unwanted background term affecting any θ_{13} measurement could indeed be useful for other applications.

Acknowledgments

I would like to thank John Bahcall, Manfred Lindner, and Carlos Peña-Garay for useful discussions and comments. This work has been supported by the W. M. Keck Foundation and the NSF grant PHY-0070928.

REFERENCES

1. S.P. Mikheev and A.Y. Smirnov, *Sov. J. Nucl. Phys.* 42 (1985) 913.
2. S.P. Mikheev and A.Y. Smirnov, *Nuovo Cim.* C9 (1986) 17.
3. L. Wolfenstein, *Phys. Rev.* D17 (1978) 2369.
4. SNO, Q.R. Ahmad et al., *Phys. Rev. Lett.* 89 (2002) 011302, *nucl-ex/0204009*.
5. KamLAND, K. Eguchi et al., *Phys. Rev. Lett.* 90 (2003) 021802, *hep-ex/0212021*.
6. M. Blennow, T. Ohlsson and H. Snellman, *Phys. Rev.* D69 (2004) 073006, *hep-ph/0311098*.
7. E.D. Carlson, *Phys. Rev.* D34 (1986) 1454.
8. Super-Kamiokande, Y. Fukuda et al., *Phys. Rev. Lett.* 82 (1999) 1810, *hep-ex/9812009*.
9. P.C. de Holanda and A.Y. Smirnov, *Phys. Rev.* D66 (2002) 113005, *hep-ph/0205241*.
10. J.N. Bahcall and C. Pena-Garay, *New J. Phys.* 6 (2004) 63, *hep-ph/0404061*.
11. C. Lunardini and A.Y. Smirnov, *Nucl. Phys.* B616 (2001) 307, *hep-ph/0106149*.
12. A.S. Dighe, M.T. Keil and G.G. Raffelt, *JCAP* 0306 (2003) 006, *hep-ph/0304150*.
13. E.K. Akhmedov, *Nucl. Phys.* B538 (1999) 25, *hep-ph/9805272*.
14. S.T. Petcov, *Phys. Lett.* B434 (1998) 321, *hep-ph/9805262*.
15. E.K. Akhmedov et al., *JHEP* 04 (2004) 078, *hep-ph/0402175*.
16. P.I. Krastev, *Nuovo Cim.* A103 (1990) 361.
17. M. Freund et al., *Nucl. Phys.* B578 (2000) 27, *hep-ph/9912457*.
18. I. Mocioiu and R. Shrock, *Phys. Rev.* D62 (2000) 053017, *hep-ph/0002149*.
19. T. Ota and J. Sato, *Phys. Rev.* D63 (2001) 093004, *hep-ph/0011234*.
20. M. Freund, P. Huber and M. Lindner, *Nucl. Phys.* B585 (2000) 105, *hep-ph/0004085*.
21. A. Cervera et al., *Nucl. Phys.* B579 (2000) 17, *hep-ph/0002108*.
22. M. Freund, *Phys. Rev.* D64 (2001) 053003, *hep-ph/0103300*.
23. P. Huber and W. Winter, *Phys. Rev.* D68 (2003) 037301, *hep-ph/0301257*.
24. P. Huber et al., *Phys. Rev.* D70 (2004) 073014, *hep-ph/0403068*.
25. P. Huber, M. Lindner and W. Winter, *Nucl. Phys.* B645 (2002) 3, *hep-ph/0204352*.
26. H. Minakata and H. Nunokawa, *JHEP* 10 (2001) 001, *hep-ph/0108085*.

27. J. Burguet-Castell et al., Nucl. Phys. B608 (2001) 301, hep-ph/0103258.
28. G.L. Fogli and E. Lisi, Phys. Rev. D54 (1996) 3667, hep-ph/9604415.
29. V. Barger, D. Marfatia and K. Whisnant, Phys. Rev. D65 (2002) 073023, hep-ph/0112119.
30. P. Huber, M. Lindner and W. Winter, Comp. Phys. Comm. 167 (2005) 195, hep-ph/0407333, <http://www.ph.tum.de/~globes>.
31. R.J. Geller and T. Hara, Phys. Rev. Lett. 49 (2001) 98, hep-ph/0111342.
32. T. Ohlsson and W. Winter, Phys. Rev. D68 (2003) 073007, hep-ph/0307178.
33. S.V. Panasyuk, REM (reference earth model) web page, <http://cfauvcs5.harvard.edu/lana/rem/index.htm>, 2000.
34. M.C. Gonzalez-Garcia and C. Peña-Garay, Phys. Lett. B527 (2002) 199, hep-ph/0111432.
35. V.D. Barger, D. Marfatia and B.P. Wood, Phys. Lett. B498 (2001) 53, hep-ph/0011251.
36. G.L. Fogli et al., Phys. Rev. D67 (2003) 093006, hep-ph/0303064.
37. J.N. Bahcall, M.C. Gonzalez-Garcia and C. Pena-Garay, JHEP 08 (2004) 016, hep-ph/0406294.
38. A. Bandyopadhyay et al., Phys. Lett. B608 (2005) 115, hep-ph/0406328.
39. M. Maltoni et al., New J. Phys. 6 (2004) 122, hep-ph/0405172.
40. CHOOZ, M. Apollonio et al., Phys. Lett. B466 (1999) 415, hep-ex/9907037.
41. G. Fogli and E. Lisi, New J. Phys. 6 (2004) 139.
42. R. Gandhi et al., (2004), hep-ph/0411252.
43. M.V. Diwan, (2004), hep-ex/0407047.